arXiv:hep-ph/0509318v1 28 Sep 2005

International Journal of Modern Physics A
© World Scientific Publishing Company

DYNAMICAL STUDY OF $\gamma^* N \leftrightarrow \Delta$ TRANSITION WITH $N(e, e'p)\pi$

SHIN NAN YANG
Department of Physics, National Taiwan University, Taipei 10617, Taiwan

SABIT S. KAMALOV
Laboratory of Theoretical Physics, JINR Dubna,141980 Moscow region, Russia

Received (Day Month Year)
Revised (Day Month Year)

The Dubna-Mainz-Taipei (DMT) dynamical model for pion electromagnetic production, which describe well the existing data from threshold up to 1 GeV photon lab energy is presented and used to analyze recent data in the Δ region . We find that within DMT model , the bare Δ is almost spherical while the physical Δ is oblate. The deformation is almost saturated by the pion cloud effects.

Keywords: Pion; Photon; Δ-resonance; $N-\Delta$ transition.

It has been well recognized that the study of the excitations of the hadrons can shed light on the nonperturbative aspects of QCD. One case which has recently been under intensive study is the electromagnetic (e.m.) excitation of the $\Delta(1232)$ resonance. At low four-momentum transfer squared Q^2, the interest is motivated by the possibility of observing a $D-$state in the Δ. The existence of a $D-$state in the Δ has the consequence that the Δ is deformed and the photon can excite a nucleon through electric $E2$ and Coulomb $C2$ quardrupole transitions, while in a symmetric SU(6) quark model the electromagnetic excitation of the Δ could proceed only via $M1$ transition. In pion electroproduction, $E2$ and $C2$ excitations would give rise to nonvanishing $E_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}$ multipole amplitudes. Recent experiments give nonvanishing ratio $R_{EM} = E_{1+}^{(3/2)}/M_{1+}^{(3/2)} \sim -0.03$ [1] at $Q^2 = 0$ which has been widely taken as an indication of the Δ deformation.

At sufficiently large Q^2, the perturbative QCD (pQCD) is expected to work. It predicts that only helicity-conserving amplitudes contribute at high Q^2, leading to $R_{EM} = E_{1+}^{(3/2)}/M_{1+}^{(3/2)} \rightarrow 1$ and $R_{SM} = S_{1+}^{(3/2)}/M_{1+}^{(3/2)} \rightarrow$ *const*. This behavior in the perturbative domain is very different from that in the nonperturbative one. It is an intriguing question to find the region of Q^2 which signals the onset of the pQCD.

In the recent measurements [2,3,4], the e.m. excitation of the Δ was studied at $0.1 < Q^2 < 4.0\ GeV^2$ via the reaction $p(e, e'p)\pi^0$. The extracted ratios R_{EM} and R_{SM} remain small and *negative*. This indicates that pQCD is still not applicable in this region of Q^2. In this talk, we want to show that the recent data of Refs. [2,3,4]

can be understood from the dominance of the pion cloud contribution at low Q^2 in the DMT dynamical model [5] for e.m. production of pion.

The main feature of the dynamical approach to the pion photo- and electroproduction [6] is that the unitarity is built in by explicitly including the final state πN interaction in the theory, namely, t-matrix is expressed as

$$t_{\gamma\pi}(E) = v_{\gamma\pi} + v_{\gamma\pi}\, g_0(E)\, t_{\pi N}(E)\,, \tag{1}$$

where $v_{\gamma\pi}$ is the transition potential operator for $\gamma^* N \to \pi N$ and, $t_{\pi N}$ and g_0 denote the πN t-matrix and free propagator, respectively, with $E \equiv W$ the total energy in the CM frame.

In the (3,3) channel the transition potential $v_{\gamma\pi}$ consists of two terms

$$v_{\gamma\pi}(E) = v^B_{\gamma\pi} + v^\Delta_{\gamma\pi}(E)\,, \tag{2}$$

where $v^B_{\gamma\pi}$ is the background transition potential which includes Born terms and vector mesons exchange contributions, as described in Ref. [7]. The second term in Eq. (2) corresponds to the contribution of a bare Δ.

We decompose Eq. (1) in the following way,

$$t_{\gamma\pi}(E) = t^B_{\gamma\pi} + t^\Delta_{\gamma\pi}(E)\,, \tag{3}$$

where

$$t^B_{\gamma\pi}(E) = v^B_{\gamma\pi} + v^B_{\gamma\pi}\, g_0(E)\, t_{\pi N}(E)\,, \qquad t^\Delta_{\gamma\pi}(E) = v^\Delta_{\gamma\pi} + v^\Delta_{\gamma\pi}\, g_0(E)\, t_{\pi N}(E)\,. \tag{4}$$

The advantage of such a decomposition is that all the processes which start with the electromagnetic excitation of the bare Δ are summed up in $t^\Delta_{\gamma\pi}$.

We evaluate $t^B_{\gamma\pi}$ with pion scattering matrix $t_{\pi N}$ obtained in a meson-exchange model [8]. Note that to make principal value integration associated with $v^B_{\gamma\pi}$ convergent, we introduce a dipole form factor with cut-off parameter Λ=440 MeV. The gauge invariance, violated due to the off-shell rescattering effects, is restored by the substitution $J^B_\mu \to J^B_\mu - k_\mu k\cdot J^B/k^2$, where J^B_μ is the electromagnetic current corresponding to the background contribution $v^B_{\gamma\pi}$ and k is the photon momentum.

For the the Δ resonance contribution $t^\Delta_{\gamma\pi}$, the corresponding multipole amplitudes A^Δ is evaluated using a Breit-Wigner form, as was done in the isobar model of Ref. [7],

$$A^\Delta(W,Q^2) \,=\, \bar{\mathcal{A}}^\Delta(Q^2)\, \frac{f_{\gamma\Delta}\, \Gamma_\Delta\, M_\Delta\, f_{\pi\Delta}}{M^2_\Delta - W^2 - iM_\Delta\Gamma_\Delta}\, e^{i\phi}\,, \tag{5}$$

where $f_{\pi\Delta}(W)$ is the usual Breit-Wigner factor describing the decay of the Δ resonance with total width $\Gamma_\Delta(W)$ and physical mass M_Δ=1232 MeV. The expressions for $f_{\gamma\Delta}$, $f_{\pi\Delta}$ and Γ_Δ are taken from Ref. [7]. The phase $\phi(W)$ in Eq. (5) is to adjust the phase of A^Δ to be equal to the corresponding pion-nucleon scattering phase $\delta^{(33)}$. Note that at the resonance energy, $\phi(M_\Delta) = 0$.

The main parameters in the bare $\gamma^* N\Delta$ vertex are the $\bar{\mathcal{A}}^\Delta$'s in Eq. (5). For the magnetic dipole $\bar{\mathcal{M}}^\Delta$ and electric quadrupole $\bar{\mathcal{E}}^\Delta$ transitions they are related to



Fig. 1. Real and imaginary parts of the, **(a)** $M_{1+}^{(3/2)}$, and **(b)** $E_{1+}^{(3/2)}$ multipoles. The open and full circles are the results from the Mainz dispersion analysis [9] and from the VPI analysis [10]. Notations for the curves are given in the text.

the conventional electromagnetic helicity amplitudes $A_{1/2}^{\Delta}$ and $A_{3/2}^{\Delta}$ by $\bar{\mathcal{M}}^{\Delta}(Q^2) = -\frac{1}{2}(A_{1/2}^{\Delta} + \sqrt{3}A_{3/2}^{\Delta})$ and $\bar{\mathcal{E}}^{\Delta}(Q^2) = \frac{1}{2}(-A_{1/2}^{\Delta} + \frac{1}{\sqrt{3}}A_{3/2}^{\Delta})$.

At the photon point $Q^2 = 0$, the bare amplitudes $\bar{\mathcal{M}}^{\Delta}(0)$ and $\bar{\mathcal{E}}^{\Delta}(0)$ of Eq. (5) were extracted from the best fit to the results of the recent analyses of Mainz [9] and VPI group [10] as shown in Fig. 1 by solid curves. The dashed curves denote the contribution from $t_{\gamma\pi}^{B}$ only. The dotted curves represented the K-matrix approximation to $t_{\gamma\pi}^{B}$, namely, without the principal value integral term included. The numerical values for $\bar{\mathcal{M}}^{\Delta}$ and $\bar{\mathcal{E}}^{\Delta}$, at $Q^2 = 0$, are given in Table. 1. Here we also give "dressed" values obtained using K-matrix approximation. One notices that the above determined bare values for the helicity amplitudes, which amount to only about 60% of the corresponding dressed values, are close to the predictions of the constituent quark model (CQM). The large reduction of the helicity amplitudes from the dressed to the bares ones results from the fact that the principal value integral part of $t_{\gamma\pi}^{B}$, which represents the effects of the off-shell pion rescattering, contributes approximately for half of the M_{1+} as indicated by the dashed curves in Fig. 1.

The Q^2 dependence of the $\bar{A}^{\Delta}$'s is parametrized by some functional form with parameters determined from fits to the data of Refs. [2,3,4]. The resulting model is found to give an excellent description on the data of Ref. [2,3,4] as well as the e.m. π^0 production near threshold [11].

Amplitudes	"bare"	"dressed"	PDG
$\bar{\mathcal{M}}^{\Delta}$	158 ± 2	289 ± 2	293 ± 8
$\bar{\mathcal{E}}^{\Delta}$	0.4 ± 0.3	-7 ± 0.4	-4.5 ± 4.2

Our extracted values for R_{EM} and R_{SM} and a comparison with the recent results of Refs. [2,3,4] are shown in Fig. 2. Our values of R_{EM} show a clear tendency to cross zero and change sign as Q^2 increases. This is in contrast with the results obtained in Refs.[2,12] which concluded that R_{EM} would stay negative with increasing Q^2.



Fig. 2. The Q^2 dependence of the ratios $R_{EM}^{(p\pi^0)}$ and $R_{SM}^{(p\pi^0)}$ at $W = 1232$ MeV. The solid and dashed curves are the DMT and MAID results, respectively. Results of our analysis obtained using MAID and experimental data from Refs.[2,3,4] (●). Open circles are from Ref.[4].

Furthermore, we find that the absolute value of R_{SM} is strongly increasing.

In summary, we calculate the Q^2 dependence of the ratios E_{1+}/M_{1+} and S_{1+}/M_{1+} in the $\gamma^* N \rightarrow \Delta$ transition, with the use of a dynamical model for electromagnetic production of pion. We find that the ratio E_{1+}/M_{1+} change the sign at $Q^2 \geq 3.0\ GeV^2$. Our results agree well with the recent measurements from Refs. [2,3,4], but deviate strongly from the predictions of pQCD. Our results indicate that the bare Δ is almost spherical and hence very difficult to be directly excited via electric E2 and Coulomb C2 quardrupole excitations. The experimentally observed $E_{1+}^{(3/2)}$ and $S_{1+}^{(3/2)}$ multipoles are, to a very large extent, saturated by the contribution from pion cloud, i.e., pion rescattering effects.

References

1. R. Beck *et al.* Phys. Rev. Lett. **78**, 606 (1997); G. Blanpied *et al.* Phys. Rev. Lett. **79**, 4337 (1997).
2. V.V. Frolov *et al.*, Phys. Rev. Lett. **82**, 45 (1999).
3. C. Mertz *et al.*, Phys. Rev. Lett. **86**, 2963 (2001)
4. K. Joo *et al.*, Phys. Rev. Lett. **88**, 122001 (2001)
5. S.S. Kamalov and S.N. Yang, Phys. Rev. Lett. **83**, 4494 (1999); S.S. Kamalov *et al.*, Phys. Rev. **C64**, 032201(R) (2001).
6. S.N. Yang, J. Phys. **G11**, L205 (1985).
7. D. Drechsel, O. Hanstein, S.S. Kamalov and L. Tiator, Nucl. Phys. **A645**, 145 (1999).
8. C.T. Hung, S.N. Yang and T.-S.H. Lee, Phys. Rev. **C64**, 034309 (2001).
9. O. Hanstein, D. Drechsel, and L. Tiator, Nucl. Phys. **A632**, 561 (1998).
10. R.A. Arndt *et al.*, Phys. Rev. **C53**, 430 (1996) (SP97 solution of the VPI analysis).
11. S.S. Kamalov *et al.*, Phys. Lett. **B522**, 27 (2001).
12. T. Sato and T.S.H. Lee, Phys. Rev. **C63**, 055201 (2001)
(2002).